

Galactic-Scale Outflow and Supersonic Ram-Pressure Stripping in the Virgo Cluster Galaxy NGC 4388

Sylvain Veilleux^{1,2}, Jonathan Bland-Hawthorn³, Gerald Cecil^{4,5}, R. Brent Tully⁶,
and Scott T. Miller¹

ABSTRACT

The Hawaii Imaging Fabry-Perot Interferometer (HIFI) on the University of Hawaii 2.2m telescope was used to map the H α and [O III] λ 5007 emission-line profiles across the entire disk of the edge-on Sb galaxy NGC 4388. We confirm a rich complex of highly ionized gas that extends ~ 4 kpc above the disk of this galaxy. Low-ionization gas associated with star formation is also present in the disk. Evidence for bar streaming is detected in the disk component and is discussed in a companion paper (Veilleux, Bland-Hawthorn, & Cecil 1999; hereafter VBC). Non-rotational blueshifted velocities of $50 - 250 \text{ km s}^{-1}$ are measured in the extraplanar gas north-east of the nucleus. The brighter features in this complex tend to have more blueshifted velocities. A redshifted cloud is also detected 2 kpc south-west of the nucleus. The velocity field of the extraplanar gas of NGC 4388 appears to be unaffected by the inferred supersonic (Mach number $M \approx 3$) motion of this galaxy through the ICM of the Virgo cluster. We argue that this is because the galaxy and the high- $|z|$ gas lie behind a Mach cone with opening angle $\sim 80^\circ$. The shocked ICM that flows near the galaxy has a velocity of $\sim 500 \text{ km s}^{-1}$ and exerts insufficient ram pressure on the extraplanar gas to perturb its kinematics. We consider several explanations of the velocity field of the extraplanar gas. Velocities, especially blueshifted velocities on the N side of the galaxy, are best explained as a bipolar outflow which is tilted by $>12^\circ$ from the normal to the disk. The observed offset between the extraplanar gas and the radio structure may be due to buoyancy or refractive bending by density gradients in the halo gas. Velocity substructure in the outflowing gas also suggests an interaction with ambient halo gas.

¹Department of Astronomy, University of Maryland, College Park, MD 20742; E-mail: veilleux@astro.umd.edu

²Cottrell Scholar of the Research Corporation

³Anglo-Australian Observatory, P.O. Box 296, Epping, NSW 2121, Australia; E-mail: jbh@aaoepp2.aao.gov.au

⁴Department of Physics and Astronomy, University of North Carolina, CB# 3255, Chapel Hill, NC 27599-3255; E-mail: cecil@noao.edu

⁵SOAR Telescope Project, NOAO, 950 N. Cherry Ave., Tucson, AZ 85726-6732

⁶Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822; E-mail: tully@galileo.ifa.hawaii.edu

Subject headings: galaxies: clusters: individual (Virgo) — galaxies: individual (NGC 4388) — galaxies: intergalactic medium — galaxies: ISM — galaxies: kinematics and dynamics – galaxies: Seyfert

1. Introduction

The edge-on Sb galaxy NGC 4388 is projected $<1^\circ$ from the core of the Virgo cluster (Biggelli, Tammann, & Sandage 1987; Yasuda, Fukugita, & Okamura 1997), but its systemic velocity of $\sim 2,525 \text{ km s}^{-1}$ is $\sim 1,500 \text{ km s}^{-1}$ above that of the Virgo cluster (e.g., Rubin, Kenney, & Young 1997; Yasuda et al. 1997). Nevertheless, studies by Chamaraux et al. (1980), Giraud (1986), Kenney & Young (1986, 1989), and Cayatte et al. (1990, 1994) show that the disk of NGC 4388 is deficient in HI; it appears truncated at $\sim 10 \text{ kpc}$, well inside the optical radius. This suggests that NGC 4388 is experiencing the effects of ram-pressure stripping by the densest portions of the intracluster medium (ICM) in the Virgo cluster. Distance estimates using the Tully-Fisher relation suggest that NGC 4388 lies close to the mean distance of the M87 subcluster in the Virgo cluster, i.e. near the core of the Virgo cluster (Mould, Aaronson, & Huchra 1980; Sandage & Tammann 1984; Pierce & Tully 1988; Yasuda et al. 1997). In the rest of this paper, we will assume that NGC 4388 is indeed a member of the Virgo cluster and will adopt a distance of 16.7 Mpc (Yasuda et al. 1997), yielding a scale of $81 \text{ pc}''$.

NGC 4388 was the first Seyfert galaxy discovered in the Virgo cluster (Phillips & Malin 1982). Nuclear activity has been detected at nearly all wavelengths. The morphology of the nuclear radio source suggests a collimated AGN-driven outflow (Stone et al. 1988; Hummel & Saikia 1991; Kukula et al. 1995; Falcke, Wilson, & Simpson 1998). The nucleus hosts a strong hard X-ray source with power-law photon index $\Gamma = 1.5$, that is heavily absorbed by a column density $N_H = 4 \times 10^{23} \text{ cm}^{-2}$ (Hanson et al. 1990; Takano & Koyama 1991; Lebrun et al. 1992; Iwasawa et al. 1997). The large equivalent width of the iron $K\alpha$ line in the ASCA spectrum of the nucleus (Iwasawa et al. 1997) and the detection of broad recombination-line emission outside (Shields & Filippenko 1988, 1996) but not at the nucleus (e.g., Miller 1992; Goodrich, Veilleux, & Hill 1994; Veilleux, Goodrich, & Hill 1997) are both consistent with a powerful hidden AGN in NGC 4388.

At optical wavelengths, NGC 4388 has been known for some time to present extended line emission (e.g., Ford et al. 1971; Sandage 1978; Colina et al. 1987; Pogge 1988; Corbin, Baldwin, & Wilson 1988). A rich complex of ionized gas extends both along the disk of the galaxy and up to $50''$ (4 kpc) above that plane. The extraplanar gas component has considerably higher ionization than the disk gas, and appears roughly distributed into two opposed radiation cones that emanate from the nucleus (Pogge 1988; Falcke et al. 1998). Detailed spectroscopic studies strongly suggest that this component is mainly ionized by photons from the nuclear continuum (Pogge 1988; Colina 1992; Petitjean & Duret 1993).

Several origins of this extraplanar gas component have been suggested: (1) it is tidal debris

from a past interaction with another galaxy (Pogge 1988); (2) supernovae have ejected it into the halo of the galaxy (e.g., Corbin et al. 1988); (3) ram pressure from the ICM has elevated some of it from the disk (Petit-Jean & Durret 1993); (4) it has been ejected from the nucleus by the mechanism that is powering the radio jet (e.g., Corbin et al. 1988).

This ambiguity stems from the weak constraints on the kinematics of the extraplanar gas that arise from the poor spatial coverage of the spectra used in previous studies. In this paper we use a Fabry-Perot spectrometer to provide complete two-dimensional coverage of the kinematics of the line-emitting region in NGC 4388. After a brief description of the data analysis methods (§2), we present the distribution and velocity field of the $H\alpha$ and [O III] $\lambda 5007$ line-emitting gas derived from the data (§3). These data allow us to split the line-emitting gas into two distinct kinematic components, one associated with the galaxy disk and the other associated with the extraplanar gas. In §4 we derive dynamical parameters of the extraplanar gas and use them to constrain the origin of this component. The important issue of ram-pressure stripping by the ICM is addressed in the context of our results. In this section, we sketch a picture of NGC 4388 that accounts for the effects of both the supersonic ICM and nuclear activity (Fig. 5). Our conclusions are summarized in §5.

2. Observations and Data Reduction

The Hawaii Imaging Fabry-Perot Interferometer (HIFI; Bland & Tully 1989) was attached at the Cassegrain focus of the University of Hawaii 2.2 m telescope at the Mauna Kea Observatory to obtain in photometric conditions 54 seven-minute exposures of redshifted $H\alpha$ and 38 eleven-minute exposures of redshifted [O III] $\lambda 5007$. The $H\alpha$ spectra were taken on 1990 May 31 and 1990 June 1 with a thick epitaxial GEC CCD ($5 e^-$ rms readout), resulting in a pixel scale of $0''.66 \text{ pixel}^{-1}$. A finesse 70 etalon with free spectral range $57 \text{ \AA}$ produced a velocity resolution of 40 km s^{-1} . The [O III] spectra were acquired on 1992 March 12 with a blue-sensitive 1024×1024 Tektronix CCD and a finesse 60 etalon having a free spectral range of $61 \text{ \AA}$. The spatial scale and spectral resolution of these data are $0''.85 \text{ pixel}^{-1}$ and 60 km s^{-1} , respectively. Order separating filters with a flat-topped transmission profile centered at $6614 \text{ \AA}$ ($55 \text{ \AA}$ FWHM) and $5030 \text{ \AA}$ ($36 \text{ \AA}$ FWHM) passed only one etalon order so true emission line profiles of $H\alpha$ and [O III] $\lambda 5007$ could be synthesized. Wavelength calibration exposures were made of diffuse Neon ($H\alpha$ cube) and Mercury-Cadmium ([O III] cube) sources. A current-stabilized, flat continuum source served to flatfield the images. Finally, the photometric standard stars HD 94808, η Hya, and θ Vir were observed through the same setup as the object to provide an absolute flux calibration.

These data were reduced using the methods described in Bland & Tully (1989), Veilleux et al. (1994), and Bland-Hawthorn (1995). However, the $H\alpha$ data suffered from several cosmetic defects associated with the aging GEC CCD. While only a fixed bias offset needed to be subtracted from the [O III] data, row-by-row bias subtraction was done on each $H\alpha$ frame to remove vertical striations of the bias pattern. After interpolating over the columns with poor charge transfer, the

images of each night were stacked in a cube in increasing order of wavelength. Each data set was flatfielded using a similarly constructed whitelight cube. This method produced good flat fields for the [O III] spectra. Large-scale illumination gradients in the sky background that persisted in the corrected H α cube were removed by dividing the spectra by a spatially smoothed sky frame. This frame was obtained by summing all of the frames in the data while interpolating through regions with H α emission from the galaxy. The photometric accuracy resulting from this method is discussed in §3.1. The rest of the reductions was the same for both H α and [O III] cubes: spatial registration of the frames, cosmic ray removal, correction for air mass variations, sky continuum and line subtraction, correction for the curvature of the isovelocity surfaces (“phase calibration”), and flux calibration. The OH sky line at 6604.13 Å (Osterbrock & Martel 1992) was also used for the velocity calibration of the H α data.

For the sake of simplicity, the observed emission-line profiles were approximated by simple Gaussian functions (regions where the emission-line profiles differ from a simple Gaussian will be discussed in §3.2). The best fits were determined by the least-chi-squared method on spectrally smoothed emission line profiles using a $1/4 - 1/2 - 1/4$ spectral filter (Hanning smoothing). Gaussian smoothing of each image with $\sigma = 1.0$ pixel improved the sensitivity to fainter features. In the end, the centroid, intensity, and width were successfully derived from $\sim 5,000$ H α spectra and $\sim 7,000$ [O III] $\lambda 5007$ spectra.

3. Empirical Results

3.1. Distribution and Excitation Properties of the Ionized Gas

Figure 1 presents the distributions of H α and [O III] $\lambda 5007$ line emission that we derived from the Gaussian fits; the underlying starlight has been removed. The $1\text{-}\sigma$ detection limit of our data corresponds to a surface brightness of ~ 7 and 3×10^{-17} erg cm $^{-2}$ s $^{-1}$ arcsec $^{-2}$ for H α and [O III] $\lambda 5007$, respectively. The flux calibration of our data cubes was checked against published long-slit spectrophotometry (e.g., Pogge 1988; and Petitjean & Durret 1993). The H α and [O III] fluxes agree with published ones to within $\sim 25 - 50\%$. Note that the H α and [O III] fluxes were not corrected for Galactic reddening because extinction is negligible toward NGC 4388 [$E(B-V) = 0.028$; Burstein & Heiles 1978, 1984]. The patchy dust lanes running roughly east-west in the optical continuum images of this galaxy (e.g., Phillips & Malin 1982) suggest highly variable internal extinction which may affect the apparent distribution of the ionized gas near the galactic plane. No attempt was made to correct for this effect. Dilution of the H α fluxes by underlying Balmer absorption is negligible in NGC 4388. Moreover, the Balmer absorption profile is dominated by pressure broadening in the atmospheres of early-type stars, making this feature indistinguishable from a slowly varying continuum over the narrow spectral bandpass of our instrument.

The most prominent features in the H α image are the strings of H II regions in the shape of

two ‘spiral arms’ within the disk of the galaxy. Strong $H\alpha$ disk emission is detected out to ~ 5 kpc east of the nucleus (its extent is limited by the velocity coverage of our data cube). The bright H II region complexes in this part of the disk each have $L_{H\alpha} \approx 10^{38.5} - 10^{39.7}$ erg s $^{-1}$. Emission near the detection limit of our $H\alpha$ data is observed projected north of the plane of the disk along $PA = 20^\circ - 50^\circ$ and 1 – 4 kpc from the nucleus. In contrast, the [O III] map is dominated by nuclear line emission. Emission from the spiral arms and from gas above the galaxy disk is also visible but considerably fainter. Its large projected $|z|$ distance implies that the gas is well above the plane of this highly inclined ($|i| = 78^\circ \pm 1^\circ$; cf. §3.2) galaxy. If this gas was in the disk, it would lie ~ 20 kpc from the nucleus which greatly exceeds the visible radius of the galaxy ($R_{25} = 14$ kpc from de Vaucouleurs et al. 1991).

The differences between the $H\alpha$ and [O III] line emission distributions is particularly evident in Figure 2, which maps the [O III]/ $H\alpha$ flux ratios. This map fundamentally compares the ionization level of the line-emitting gas. Energies of 35.5 eV and 54.9 eV are needed to produce O^{++} and O^{+++} , i.e. significantly larger than the 13.6 eV needed to ionize hydrogen. Uncertainties on ratios range from 25% in the bright H II regions in the disk to $>50\%$ in the fainter high- $|z|$ gas (recall that the extraplanar $H\alpha$ emission is near the detection limit of the data). The low-ionization gas is found in the bright H II regions of the spiral arms. These H II regions have [O III] $\lambda 5007/H\alpha$ ratios of ~ 0.1 , typical of star-forming regions with low ionization temperature ($\sim 40,000$ K; Evans & Dopita 1985) or high metal abundance ($O/H \sim 3 \times O/H_\odot$; McCall, Rybski, & Shields 1985). The high-ionization gas concentrates into two regions north and south of the nucleus. The southern extension has a conical shape with apex near the nucleus. Typical values for the [O III] $\lambda 5007/H\alpha$ ratio in the nucleus and in the ionization cones are 1 to 5, consistent with photoionization by the AGN (Pogge 1988; Colina 1992; Petitjean & Durret 1993). This ionization map is qualitatively similar to previously published [O III] $\lambda 5007/(H\alpha + [N II] \lambda\lambda 6548, 6583)$ ratio maps obtained with narrow-band interference filters (Pogge 1988; Corbin et al. 1988; Wilson & Tsvetanov 1994; Mulchaey, Wilson, & Tsvetanov 1996; Falcke et al. 1998). Note, however, that the present line ratio map was derived directly from our spectra and is therefore not subject to the large uncertainties in the line separation and continuum subtraction which often affect line ratio maps derived from narrow-band images.

3.2. Kinematics of the Extraplanar Material

Figure 3 presents the barycentric velocity fields that are derived from the $H\alpha$ and [O III] $\lambda 5007$ spectra; uncertainties range from ~ 20 km s $^{-1}$ in the bright line-emitting regions to $\gtrsim 100$ km s $^{-1}$ in the fainter areas. The overall velocity field of the [O III]-emitting gas resembles that of the $H\alpha$ -emitting gas. In both, the velocity field in the disk is characterized by a large-scale east-west gradient indicative of rotation in the plane of the disk. The rotation curve derived from our data is highly consistent with earlier results (cf. Rubin, Kenney, & Young 1997 and references therein). Non-Gaussian $H\alpha$ and [O III] $\lambda 5007$ profiles are found near the nucleus ($R \lesssim 10''$), again

confirming earlier results (Iye & Ulrich 1986; Colina et al. 1987; Ayani & Iye 1989; Veilleux 1991; Rubin et al. 1997).

Figure 3 shows a kinematic dichotomy between the disk gas and the extraplanar gas. To more objectively identify which portion of the line-emitting gas does not partake in disk rotation, we have constructed kinematic models to simulate the observed velocity field in the disk. These models are described in more detail in VBC. Because the focus of the present paper is on the origin of the extraplanar gas in NGC 4388, only the main results of our analysis of the disk velocity field will be discussed here.

To minimize possible effects of the active nucleus, our analysis of the disk component focused on the $H\alpha$ velocity field. To summarize our results, Figure 4 compares the observed $H\alpha$ velocity field with our best-fitting axisymmetric disk model and our best-fitting disk model that includes bar streaming motions. Compared to the model with purely circular motions, the velocity residuals from the model incorporating elliptical streaming are significantly smaller throughout the disk. The residuals have a symmetric Gaussian distribution with dispersion of only 10 km s^{-1} . This is an excellent fit considering that hydrodynamic effects (e.g., strong shocks along the bar; Athanassoula 1992; Piner, Stone, & Teuben 1995) are ignored in our simple kinematic model. The parameters of the model with elliptical streaming are listed in Table 1 and discussed in more detail in VBC. The strong bar streaming motions prevented us from deriving a rotation curve directly from the data; instead, we used a smooth parametric function to model the rotation curve (cf. eqn. 3 in VBC). Note that the inclination of the disk, $i = -78^\circ$, implies that the north rim of the disk is the near side (i.e. we are looking at the galaxy from under the disk).

Our kinematic model suggests that most of the gas in the disk of NGC 4388 partakes in bar-forced orbital motion around the nucleus. There are, however, two notable exceptions. The $H\alpha$ and [O III]-emitting gas with $R \gtrsim 1.2 \text{ kpc}$ and $PA = 10^\circ - 70^\circ$ has velocities which appear blueshifted by $75 - 250 \text{ km s}^{-1}$ relative to systemic velocity. The location and velocity field of this gas implies that it is elevated above the disk of the galaxy.

No obvious kinematic trends with height above the disk or with galactocentric radius are seen in the NE complex. However, a correlation appears to exist between the velocity field and the surface brightness of the gas in the NE complex. The most extreme velocities in this gas complex appear in the bright cloud located near the nucleus at $R \approx 2 \text{ kpc}$ and $PA \approx 55^\circ$ ($v = -250 \text{ km s}^{-1}$; the ‘NE cloud’ in the nomenclature of Pogge 1988). Double-peaked [O III] profiles with velocity splitting $\sim 140 \text{ km s}^{-1}$ are seen near the north-east edge of that cloud. The much fainter gas directly east of this NE cloud has velocities which average 175 km s^{-1} larger (-75 km s^{-1} relative to systemic) than that of the cloud. The trend between surface brightness and blueshifted velocities is also seen in the cloud complex at $R \approx 3.3 \text{ kpc}$ and $PA \approx 35^\circ$. Here, the brighter clouds have velocities which are $\sim 60 - 100 \text{ km s}^{-1}$ more blueshifted than the diffuse gas surrounding them.

The bright [O III] cloud south-west of the nucleus at $R \approx 1.8 \text{ kpc}$ and $PA = 215^\circ$ also

has anomalous velocities relative to the disk. This cloud is not detected at $H\alpha$, implying $[O\ III]\ \lambda 5007/H\alpha \gtrsim 5$. Substructures (shoulders, line splitting) of order $\sim 200\ \text{km s}^{-1}$ are seen in the emission-line profiles on the north-east side of that cloud (also detected by Corbin et al. 1988). A comparison with the modeled disk velocity field indicates that this cloud has an average redshift of $\sim 100 - 150\ \text{km s}^{-1}$ relative to the underlying disk gas. Such supersonic motion through the disk ISM is highly unlikely; therefore, this cloud is probably in front of the disk. Its radial velocity relative to systemic is only $\sim 20 - 30\ \text{km s}^{-1}$.

Contrary to Petitjean & Durret (1993), we see no general trend for gas at large radii to have negative velocities relative to the galaxy rotation. The probable origin of this discrepancy is the use of different kinematic models to reproduce the observed velocity field. As we demonstrated in §3.1, elliptical streaming is important in the disk of NGC 4388 but this effect was not incorporated in the axisymmetric models of Petitjean & Durret (1993). We feel that the complete two-dimensional coverage of our spectra and our use of slightly more sophisticated models put our conclusions on a more solid basis.

4. Discussion

4.1. Derived Properties of the Extraplanar Material

A clue to the origin of the extraplanar gas is provided by the energetics of this material. In this section, we estimate the dynamical time scale, mass, and kinetic energy of the extraplanar gas in NGC 4388.

An upper limit on the dynamical time scale of the extraplanar gas can be estimated from its observed velocity and linear extent: $t_{\text{dyn}} \approx R/V \lesssim 2 \times 10^7\ (R/4\ \text{kpc})\ (V_{\text{obs}}/200\ \text{km s}^{-1})^{-1}\ \text{yr}$. The ionized mass above the disk follows from the $[O\ III]\ \lambda 5007$ flux. Other researchers have estimated the density and temperature of the extraplanar gas. Pogge used the $[S\ II]\ \lambda 6731/\lambda 6716$ flux ratio to find densities in the range $100 - 1,000\ \text{cm}^{-3}$. However, densities $< 50\ \text{cm}^{-3}$ are claimed by Colina (1992) and Petitjean & Durret (1993) for the NE complex. Estimates for the temperature range from $\sim 10,000\ \text{K}$ (Petitjean & Durret 1993) to $15,000\ \text{K}$ (Colina 1992). In the following discussion, we parametrize the mass in terms of the density and use an electron temperature of $10^4\ \text{K}$.

The total $[O\ III]$ intensity of $\sim 2.5 \times 10^{-13}\ \text{erg s}^{-1}\ \text{cm}^{-2}$ beyond $R = 1.2\ \text{kpc}$ and within $PA = 10 - 70^\circ$ yields an ionized mass in the NE complex of $M(\text{NE}) \approx 4 \times 10^4\ X^{-1}\ N_{e,2}^{-1}\ M_\odot$, where X is the fraction of doubly ionized oxygen and $N_{e,2}$ is the electron density in units of $100\ \text{cm}^{-3}$. In this calculation, we used a solar mass fraction for oxygen and a five-level atom approximation to estimate the emission coefficient (McCall 1984). A lower limit on the bulk kinetic energy of the north-east gas, $E_{\text{bulk}}(\text{NE})$, may be derived by summing the bulk kinetic energy over each pixel in this region ($\sum_i 1/2\ \delta m_i\ v_i^2$, where δm_i and v_i are the mass and observed radial velocity at pixel

i). We find $E_{\text{bulk}}(\text{NE}) \gtrsim 6 \times 10^{51} X^{-1} N_{e,2}^{-1}$ ergs. The “turbulent” kinematic energy of this gas, $E_{\text{turb}}(\text{NE})$, may be derived from $\Sigma_i \delta m_i \sigma_i^2$, where $\sigma_i = V_{\text{FWHM}}/1.67$; we get $E_{\text{turb}}(\text{NE}) \approx 1 \times 10^{52} X^{-1} N_{e,2}^{-1}$ ergs. Following a similar procedure for the SW cloud, we find $M(\text{SW}) \approx 1,000 X^{-1} N_{e,2}^{-1} M_{\odot}$, $E_{\text{bulk}}(\text{SW}) \gtrsim 6 \times 10^{49} X^{-1} N_{e,2}^{-1}$ ergs and $E_{\text{turb}}(\text{SW}) \approx 3 \times 10^{50} X^{-1} N_{e,2}^{-1}$ ergs. Using $X = 1$ and $N_{e,2} = 0.1$, the total mass and kinetic energy of the extraplanar gas are thus $\sim 4 \times 10^5 M_{\odot}$ and $E_{\text{kin}} = E_{\text{bulk}} + E_{\text{turb}} \gtrsim 1 \times 10^{53}$ ergs. Using our estimate of the dynamical timescale, we derive a lower limit on the time-averaged rate of kinetic energy injected into the extraplanar gas: $dE_{\text{kin}}/dt \simeq E_{\text{kin}}/t_{\text{dyn}} \gtrsim 2 \times 10^{37} N_{e,2}^{-1} \text{ erg s}^{-1}$. We must emphasize that the energetics derived in this section only take into account the optical component of the extraplanar gas. Entrainment of the molecular gas from the disk could significantly increase the energetics of the extraplanar material.

4.2. Origin of the Extraplanar Material

Our results can constrain the origin of the extraplanar gas in NGC 4388. In this section, the predictions of each of the four scenarios introduced in §1 are discussed and compared with our data.

4.2.1. Tidal Debris from a Past Galaxy Interaction

In analogy with the Magellanic Stream (Mathewson, Schwarz, and Murray 1977; Lin, Jones, & Klemola 1995), the extraplanar material in NGC 4388 may represent the tidal debris from a recent encounter with a small gas-rich galaxy (Pogge 1988). In this scenario, gas tidally stripped from the dwarf galaxy appears when ionized by the radiation cone of the active nucleus. This scenario has been suggested by Tsvetanov & Walsh (1992) to explain the narrow ionized “arcs” in NGC 5252. In this S0 galaxy with an AGN, the ionized arcs are invisible outside sharp-edged conical sectors. However, the arcs have kinematics (Morse et al. 1998) and space distribution that continue into H I gas (Prieto & Freudling 1996) on a ring inclined to the S0 disk that is biconically illuminated, hence ionized, by nuclear photons.

Numerical stellar dynamical simulations of the accretion of a dwarf galaxy by a large disk galaxy show that the satellite sinks in only ~ 1 Gyr and causes considerable damage to the disk’s inner regions (Walker, Mihos, & Hernquist 1996). Satellites on bound polar orbits experience severe orbital decay, settling into an orbit coplanar with the disk after only one or two orbits (Walker et al. 1996). The well-behaved velocity field of the disk in NGC 4388 and the presence of gas well above the disk therefore tightly constrains the epoch of the purported encounter, i.e. $\lesssim 10^8$ yrs. In this scenario, the velocity field of the tidal debris is expected to reflect the orbital motion of the original dwarf galaxy, showing trends with radial distance from the nucleus, the center of the potential well. These trends are absent in our data. Moreover, it is difficult

to explain in this scenario the velocity – surface brightness correlation and the steep velocity gradients observed in the NE complex (§3.2).

4.2.2. Interaction with the Intracluster Medium

NGC 4388 is one of the most HI-deficient spiral galaxies in the Virgo cluster (Chamaraux et al. 1980; Cayatte et al. 1990; 1994). A group III galaxy in the nomenclature of Cayatte et al. (1994), it has a severely truncated HI disk with an HI-to-optical size ratio $D_{\text{HI}}/D_{\text{opt}} = 0.7$. NGC 4388 is located very close to the center of the Virgo cluster. Taking the position of the core to be the peak position of the luminosity density, $\alpha = 12^{\text{h}} 25.3^{\text{m}}$ and $\delta = +13^{\circ} 18'$ and adopting 16.5 Mpc for the core distance (Binggeli, Tammann, & Sandage 1987; Yasuda et al. 1997), we find that NGC 4388 is <300 kpc from the center of the Virgo cluster. Using $V_{\text{Virgo}} \approx 1,000 \text{ km s}^{-1}$ for the recession velocity of the cluster (Yasuda et al. 1997), NGC 4388 is plunging edgewise at $-1,500 \text{ km s}^{-1}$ into the core. It is therefore a prime candidate for ram pressure and turbulent viscous stripping by the ICM.

Ram-pressure stripping has been proposed by Petitjean & Durret (1993) to explain the velocity field of the extraplanar material in NGC 4388. In this scenario, the line-emitting gas in the NE complex may represent (1) gas that has cooled and condensed out of the ICM or (2) interstellar gas originally in the disk of the galaxy which has been elevated by ram-pressure exerted by the fast-moving ICM. The first possibility appears unlikely on the basis that the cooling time scale of the ICM (Rangarajan et al. 1995),

$$t_{\text{cool}} \approx 4 \times 10^6 (T_{\text{ICM}}/10^7 \text{ K})^{3/2} (n_{\text{ICM}} \text{ cm}^{-3})^{-1} \text{ yr}, \quad (1)$$

exceeds the Hubble time for the conditions of the Virgo ICM ($T_{\text{ICM}} = 2.5 \text{ keV} = 2 \times 10^7 \text{ K}$, $n_{\text{ICM}} \approx 10^{-4} \text{ cm}^{-3}$; Koyama, Takano, & Tawara 1991; Böhringer et al. 1994). We therefore focus on the second scenario.

We suppose that the extraplanar gas arises from interstellar gas that has been ram-pressure stripped from the galactic disk. In this scenario, the ram pressure exerted by the ICM will be largest on the *receding* eastern edge of the disk so this is the most likely source of extraplanar gas; this is consistent with the presence of high- $|z|$ gas east of the nucleus. The swept-out gas will be heated conductively by the hot ICM; it is therefore important to determine if this gas will survive long enough in the ICM environment to be detected optically at the observed large heights above the disk. Here, we consider only the HI gas phase because the molecular gas distribution in NGC 4388 appears normal (Kenney & Young 1989). Following Cowie & McKee (1977), we calculate the evaporation timescale for a cloud embedded in the ICM. In the galaxy halo, classical thermal conduction (i.e. no ordered magnetic fields that would reduce the conductivity; see Hummel & Saikia 1991) should provide a reasonable estimate for the evaporation time, which is given by

$$t_{\text{evap}} = 1,000 n_c R_{\text{pc}}^2 (T_{\text{ICM}}/10^7 \text{ K})_{\text{ICM}}^{-5/2} (\ln \Lambda/30) \text{ yr} \quad (2)$$

where n_c is the embedded cloud density, R_{pc} is its radius in parsecs, T_{ICM} is the temperature of the surrounding hot medium, and Λ is the Coulomb logarithm. For typical Galactic HI clouds ($n_c \approx 1 \text{ cm}^{-3}$, $R \approx 10 \text{ pc}$) embedded in a hot medium of temperature 2.5 keV (Koyama, Takano, & Tawara 1991; Böhringer et al. 1994), the evaporation time is $t_{\text{evap}} \approx 10^4 \text{ yrs}$. This is much shorter than the dynamical timescale $t_{\text{dyn}} \approx 2 \times 10^7 \text{ years}$ derived in §4.1.

Shock waves generated during the sweeping may also contribute to heating the interstellar clouds. The radial velocity $V_r = 1,500 \text{ km s}^{-1}$ of NGC 4388 relative to the ICM would produce shock waves in the ICM (where the sound speed $c_{ICM} \approx 0.1 T_{ICM}^{1/2} \text{ km s}^{-1} \approx 500 \text{ km s}^{-1}$) and in the interstellar gas of NGC 4388 ($c_{HI} < 3 \text{ km s}^{-1}$, assuming HI with $T < 10^3 \text{ K}$). The shock velocity in the HI cloud will be $v_s \approx (n_{ICM}/n_c)^{1/2} V_r \approx 15 \text{ km s}^{-1}$ (e.g., McKee & Cowie 1975; Sgro 1975) and the gas in the cloud will be heated to $T \approx 14 v_s^2 \approx 3,200 \text{ K}$ in a time $t_h \sim 2R/v_s \approx 10^6 \text{ yrs} \ll t_{\text{dyn}}$.

Another serious argument against the ram-pressure hypothesis is based on the kinematics of the extraplanar gas. The equation of motion for the HI clouds that are stripped by the ICM and somehow survive evaporation/heating is

$$M_c g = C \rho_{ICM} (V_{ICM} - V_c)^2 l_c^2, \quad (3)$$

where g is the cloud acceleration, M_c , V_c , and l_c , are the mass, speed, and characteristic size of the interstellar clouds, and ρ_{ICM} and V_{ICM} are the mass density and speed of the ICM, respectively. The value of the constant C depends on the details of the cloud model (e.g., $C \approx 9$ for an isothermal cloud model; DeYoung & Axford 1969; Longair, Ryle, & Scheuer 1973; Blandford & Konigl 1979a,b). The equation of motion can be simplified if we make the reasonable assumption that $V_c \ll V_{ICM}$ in the present situation and also assume that the size and mass of the clouds are constant. (In reality the cloud expands because of the decrease in ram pressure and its mass may also decrease because of erosion caused by instabilities). With these assumptions, the distance x_{acc} over which the cloud needs to be accelerated to attain velocity V_c is given by

$$x_{\text{acc}} = \frac{V_c^2}{2g} = \left(\frac{M_c V_c^2}{2C \rho_{ICM} V_{ICM}^2 l_c^2} \right). \quad (4)$$

Using $V_c \approx 200 \text{ km s}^{-1}$ for the velocity of the extraplanar gas and $M_c \approx 20 M_\odot$ for the mass of typical HI clouds ($n_c \approx 1 \text{ cm}^{-3}$; $l_c \approx 10 \text{ pc}$) entrained in the ICM, we get x_{acc} of only $\sim 100 \text{ pc}$.

This simple model therefore predicts velocities for the extraplanar material at $\sim 2 - 4 \text{ kpc}$ which considerably exceed those observed. Large projection effects where much of the velocity of the entrained gas is in the plane of the sky are unlikely. The location and very large velocity of NGC 4388 relative to the Virgo cluster ($V_r = 1,500 \text{ km s}^{-1} \gg \sigma_{\text{Virgo}} = 762 \text{ km s}^{-1}$, the Virgo cluster velocity dispersion; Rangarajan et al. 1995) strongly suggests that NGC 4388 is on a radial orbit roughly aligned along our line of sight (otherwise, NGC 4388 would not be bound to the Virgo cluster; see also Giraud 1986). The observed tendency for gas at large heights above the disk to have velocities close to systemic is opposite to that expected for ISM entrained in the

ICM. (To first order, we expect $V_c \propto |z|^{1/2}$ in our simple model; eqn. 4). Nor can the brightness – velocity correlation in the NE complex and the redshifted velocity of the SW cloud be explained in this scenario. We conclude that the extraplanar gas component is not exposed to the putative, high velocity ICM. We discuss this surprising result in more detail in §4.3.

4.2.3. *Supernova-Driven Event*

Corbin et al. (1988) have suggested that part of the NE complex (more specifically, the NE cloud) may have been elevated above the disk by a recent supernova. This scenario deserves serious consideration because evidence for intense star formation is seen throughout the disk of NGC 4388 (Fig. 1). Associated with these star-forming complexes are clustered supernovae that deposit significant kinetic energy into the surrounding ISM. The kinetic energy from the stellar ejecta will thermalize rapidly to create an expanding “bubble” of very hot ($T \approx 10^7 - 10^8$ K) gas. Following Mac Low & McCray (1988) and Mac Low, McCray, & Norman (1989), the condition for this bubble to “break out” of the disk is given by

$$D \simeq 3 \times 10^{-42} (dE/dt) H_{kpc}^{-2} P_7^{-3/2} n_0^{1/2} > 100, \quad (5)$$

where dE/dt is the rate at which energy is injected in the disk, H_{kpc} is the scale height of the galactic disk in kpc, n_0 is the ambient density in cm^{-3} , and P_7 is the ambient pressure, P/k , in units of 10^7 K cm^{-3} . In analogy with our analysis of the disk of NGC 3079 (Veilleux, Bland-Hawthorn, & Cecil 1995), we assume that the multi-phase nature of the ISM in NGC 4388 can be simplified by using the following average disk parameters: $H_{kpc} = 1.0$, $n_0 = 1.0$, and $P_7 = 10^{-3}$. Under these circumstances, break-out occurs if $dE/dt \gtrsim 1 \times 10^{39} \text{ erg s}^{-1}$. Note that this analysis neglects cushioning by the ambient magnetic field (cf. Tomisaka 1990; Ferrière, Mac Low, & Zweibel 1991; Slavin & Cox 1992; Mineshige, Shibata, & Shapiro 1993). We have therefore probably underestimated the energy injection rate required for break-out.

Are any of the star-forming regions in the disk of NGC 4388 sufficiently powerful to eject the NE cloud out of the disk? Based on the projected location of the NE cloud, the best candidate HII region complex for this event lies at $R \approx 1.9$ kpc east of the nucleus. This complex has $L(\text{H}\alpha) \approx 2 \times 10^{39} \text{ erg s}^{-1}$. Using equation (4) of Veilleux et al. (1995) to translate this $\text{H}\alpha$ luminosity into a time-averaged rate of energy injection from stellar winds and supernovae, we obtain $dE/dt \approx 3 \times 10^{39} \text{ erg s}^{-1}$, barely sufficient for break-out. Note that our analysis has not accounted for the kinematic energy in the NE complex itself: $E_{\text{kin}}(\text{NE cloud}) = E_{\text{bulk}}(\text{NE cloud}) + E_{\text{turb}}(\text{NE cloud}) \gtrsim 5 \times 10^{51} \text{ ergs}$, is quite substantial – the equivalent to the energy released from $\gtrsim 5$ Type II SNe.

Our dynamical analysis suggests that it is difficult to explain the NE cloud as a result of star formation. But there are two important assumptions. First, we neglected dust obscuration by the edge-on disk when deriving the energy injection rate from the $\text{H}\alpha$ flux of the eastern HII region. Extinction estimates towards the nucleus ($A_V \approx 1 - 2$ mags; Phillips & Malin 1982; Storchi-Bergmann et al. 1990; Osterbrock et al. 1992; Petitjean & Durret 1993) suggest that

the correction may be substantial. Second, it is unclear if the *current* energy injection rate from this complex is relevant to an event that presumably took place $z(\text{NE})/V(\text{NE}) \approx 10^7$ years ago. Evidently, dynamical arguments alone fail to reject the supernova hypothesis.

Nonetheless, we consider this hypothesis unlikely for three further reasons: (1) Perhaps most damaging is the morphology of the NE complex. It does not resemble the filaments/loops produced by intense star formation in the disks of galaxies (e.g., Dettmar 1992; Veilleux et al. 1995). A bridge of blueshifted gas between the NE cloud and the nucleus is detected in our [O III] data. This is strongly suggestive of mass transfer from the nucleus to the NE cloud rather than from the disk (see next section). (2) The overall blueshifted velocity field of the NE complex is difficult to explain in this scenario. Gas ejected by SN-driven events is expected to have a strong velocity component perpendicular to the plane of the disk (e.g., Norman & Ikeuchi 1989). Because $i(\text{NGC 4388}) = -78^\circ$, we expect that most of this gas will be traveling *away* from us, not toward us. In addition, the SN-event hypothesis cannot explain the presence of the SW cloud. (3) The relatively long orbital period in the disk at $R \approx 2$ kpc ($2\pi R/V_{\text{orb}} \approx 10^8$ years $> t_{\text{dyn}}$) implies that morphological (e.g., filaments, bubbles) or kinematical (e.g., non-rotational motion) disturbances caused by a supernova should still be visible in the disk. None are detected in our data.

4.2.4. Nuclear Outflow

NGC 4388 has an elongated, jet-like radio structure of nonthermal origin which extends ~ 200 pc south of the nucleus (e.g., Stone et al. 1988; Carral, Turner, & Ho 1990; Hummel & Saikia 1991; Kukula et al. 1995; Falcke et al. 1998). This suggests a collimated nuclear outflow. This radio morphology resembles that of several other Seyfert galaxies where the ejected radio plasma is clearly interacting with the surrounding ISM, producing elongated optical narrow-line regions where the gas is compressed and entrained in the outflow, but is probably kept ionized by the nuclear continuum (e.g., Whittle et al. 1988).

Is this phenomenon occurring in NGC 4388? The orientation of the putative nuclear outflow in NGC 4388 does not favor strong interaction between radio plasma and ISM in the galaxy disk: the nuclear radio source (i.e. the nuclear component and “blob A” in the nomenclature of Hummel & Saikia 1991) is elongated along P.A. $\approx 23^\circ$, i.e. close to the minor axis of the galaxy disk. A diffuse bubble-like structure extends ~ 1.5 kpc above the nucleus due north (P.A. $\approx 5^\circ$) of the galaxy disk. This strongly suggests a nuclear outflow which has burst out of the galaxy plane. The existence in Figure 3 of a continuous bridge of approaching (blueshifted) gas between the nucleus and the NE complex also supports this picture. The presence of a redshifted cloud south-west of the nucleus, i.e. diametrically opposite to the blueshifted NE complex suggests that the outflow is bipolar. The mass and kinetic energy involved in the extraplanar gas (cf. §4.1) are similar to those involved in galactic-scale outflows in nearby starburst and Seyfert galaxies (e.g., Cecil, Bland, & Tully 1990; Heckman, Armus, & Miley 1990; Veilleux et al. 1994; Veilleux & Bland-Hawthorn 1997; Shopbell & Bland-Hawthorn 1998). Line splitting near the NE and SW clouds may reflect

regions of high dissipation of kinetic energy caused by the interaction/deflection of the outflowing gas with ambient halo gas (e.g., Gallimore et al. 1996a, 1996b). The blueshifted velocities of the extraplanar gas above the north (near) side of the galaxy and redshifted velocities below the south (far) side suggests that the radio jet tilts relative to the normal to the disk by at least $90^\circ - |i| \approx 12^\circ$.

One difficulty with the outflow scenario is the lack of correlation between radio features in NGC 4388 and the extraplanar optical gas. Standard plasmon-driven bowshock models predict a one-to-one correspondence between the radio and optical structures (e.g, Pedlar, Dyson, & Unger 1985; Taylor et al. 1989). While the lack of obvious optical counterpart to radio blob A can be attributed to dust obscuration by the disk, the origin of the large offset between the NE line-emitting complex in NGC 4388 and the northern radio bubble is harder to explain. The NE complex is three times larger than the northern radio bubble, and is misaligned from the axis of the radio bubble by $\sim 30^\circ$.

However, morphological differences between radio and optical features are common among edge-on galaxies with confirmed large-scale galactic outflows (e.g., Colbert et al. 1996a, 1996b, 1998; Lehnert & Heckman 1996). In NGC 3079, for instance, most of the radio emission lies outside the kpc-scale optical nuclear superbubble (Veilleux et al. 1994), and X-shaped optical filaments with no obvious radio counterparts rise more than 4 kpc above the disk plane (Heckman, Armus, & Miley 1990; Veilleux et al. 1995). Several processes may explain the lack of correlation between radio and optical emission in NGC 4388. The lighter radio plasma is more susceptible to external forces than the denser optical filaments. Buoyancy due to density differences between the radio cloud and ambient gas in the halo and the galactic gravitational gradient (parallel to the minor axis of the disk) have been suggested to explain the bending in the radio structure (Stone et al. 1988). Here the net buoyancy force on a cloud of density ρ_c immersed in a region of density ρ_0 is $\propto \rho_0 - \rho_c$. Buoyancy will have no direct effects on the denser optical line-emitting gas, thereby kinematically decoupling this gas from the radio plasma. Balancing the buoyancy on the radio cloud against the ram pressure from the ambient halo gas and assuming $\rho_c \ll \rho_0$, Stone et al. (1988) derived a terminal velocity for the radio cloud of $10 - 35 \text{ km s}^{-1}$, i.e. much smaller than the apparent velocity of the *line-emitting* gas. Interestingly, the time needed to reach the current radio extent at this speed is similar to the dynamical timescale for the line-emitting gas ($t_{\text{dyn}} \approx 2 \times 10^7 \text{ yrs}$). The lighter radio plasma will also be more subject to refractive bending by density gradients in the halo gas than the dense line-emitting gas (Henriksen, Vallée, & Bridle 1981; Smith & Norman 1981; Fiedler & Henriksen 1984). A steep vertical density gradient in the ambient gas will force the radio plasma to follow a path close to the minor axis of the galaxy, as seen in the radio data. Finally, magnetic forces may also act to confine the lighter radio plasma (e.g., Wehrle & Morris 1987) without substantially affecting the kinematics of the more massive optical filaments. Current radio data do not constrain the strength and orientation of the magnetic field in the extraplanar gas component.

Assuming the morphological differences between the radio and optical emission are indeed

due to a kinematic decoupling between these two gas components, one question still remains: is the radio jet in the nucleus of NGC 4388 powerful enough to accelerate the gas in the NE complex to its current velocity? To answer this question we assume that the acceleration phase of the line-emitting gas occurred near the nucleus when the radio plasma was still coupled to the line-emitting gas. The line-emitting clouds, initially at rest, were accelerated downstream by the jet. Under these assumptions, the equation of motion of the line-emitting clouds at this early epoch is similar to that of clouds entrained in the ICM (eqn. 3). The distance x_{acc} over which the cloud must accelerate to reach velocity V_c is given by equation (4) after replacing ρ_{ICM} and V_{ICM} by ρ_j and V_j , the mass density and velocity of the jet gas. The value of x_{acc} can be estimated if we use the fact that $L_r = \epsilon N_c \rho_j V_j^3 l_c^2 / 2$ where L_r is the luminosity of the radio jet, N_c is the number of clouds entrained by this jet, and ϵ is the fraction of the total power of the jet which is converted into radio emission (Wilson 1981):

$$\begin{aligned} x_{acc} &= \frac{\epsilon M_{tot} V_c^2 V_j}{4 C L_r}, \\ &= \left(\frac{2 \epsilon}{10^{-2}} \right) \left(\frac{M_{tot}}{10^6 M_\odot} \right) \left(\frac{V_c}{200 \text{ km s}^{-1}} \right)^2 \left(\frac{V_j}{10^4 \text{ km s}^{-1}} \right) \left(\frac{10}{C} \right) \left(\frac{10^{37} \text{ erg s}^{-1}}{L_r} \right) \text{ kpc.} \end{aligned} \quad (6)$$

Here, $M_{tot} = N_c M_c \simeq 4 \times 10^5 M_\odot$ is the total mass of line-emitting clouds entrained in the radio jet (cf. §4.1). As a measure of the jet radio luminosity, we use the sum of the radio luminosity in “blob A” and the “elongated feature” (Hummel & Saikia 1991; $L_r \approx 7 \times 10^{37} \text{ erg s}^{-1}$ for our adopted distance of 16.7 Mpc). A reasonable estimate for x_{acc} is therefore $\sim 100 \text{ pc}$, implying that the acceleration zone of the line-emitting clouds is well within the radial position of the NE cloud ($R \approx 2 \text{ kpc}$). Entrainment with the radio jet can therefore produce the observed velocities of $200 - 300 \text{ km s}^{-1}$. While the previous derivation neglects projection effects and ram pressure by the ambient halo gas (both of which will tend to increase the required acceleration distance, x_{acc}), we feel that the kinematic energy in the NE complex can be explained through this nuclear outflow scenario. From equation (3), ram-pressure deceleration by the ambient halo gas will be stronger for small clouds (large l_c^2/M_c) than for large clouds, and may therefore explain the larger blueshift of the brighter (more massive) clouds in the NE complex.

4.3. Nature of the Interaction with the Virgo Intracluster Medium

The disk of NGC 4388 is oriented ($i = -78^\circ$) such that the extraplanar gas to the north-east of the galaxy should feel the full force of the ICM ‘wind’, while the high- $|z|$ gas to the south may be somewhat ‘shielded’ by the galaxy, lying in the wake of the flow. Simulations of radio jet propagation in a moving ICM (e.g., O’Donohue, Eilek, & Owens 1993; Loken et al. 1995) show that the ram pressure from the ICM should have profound effects on the morphology and kinematics of the outflowing gas. The lack of evidence for entrainment of the north-east extraplanar gas in the Virgo ICM is therefore puzzling. In this section we discuss in more detail

the nature of the interaction between NGC 4388 and the Virgo ICM, and estimate the effects of this interaction on the kinematics of the extraplanar material.

We first re-examine the rationale for ICM–ISM interaction in NGC 4388. Ram-pressure sweeping depends on the local pressure of the intracluster medium, the relative motion of a galaxy relative to the intracluster medium and the local surface density within the galaxy. Ram-pressure stripping of the gaseous phase of a galaxy will occur if the ram pressure from the ICM exceeds the gravitational restoring force in the galaxy (Gunn & Gott 1972), i.e.

$$\rho_{\text{ICM}} V_{\perp}^2 > 2\pi G \sigma_{\text{tot}} \sigma_{\text{gas}} \quad (7)$$

where ρ_{ICM} is the density of the intracluster medium, $V_{\perp}$ is the relative velocity of the intracluster medium perpendicular to the galaxy plane, σ_{tot} is the galaxy total surface density and $\sigma_{\text{gas}} = \sigma_{\text{HI}} + \sigma_{\text{H}_2}$ is the gas surface density. Using $V_{\perp} = 1,200 \text{ km s}^{-1}$, $\rho_{\text{ICM}} = 1.6 \times 10^{-4} \text{ cm}^{-3}$ (derived from a general model fit to the X-ray surface brightness profile; Takano et al. 1989, and references therein), $\sigma_{\text{gas}} = \sigma_{\text{HI}} = 10^{20} \text{ cm}^{-2}$, and $\sigma_{\text{tot}} = 157 \text{ M}_{\odot} \text{ pc}^{-2}$ (calculated using a Brandt model potential with $n = 1.3$ at the HI radius where $\sigma_{\text{gas}} = 10^{20} \text{ cm}^{-2}$), Cayatte et al. (1994; their Table 3) have argued that condition (7) is met beyond $R \approx 6 \text{ kpc}$ in NGC 4388. But the large value of $V_{\perp}$ used in their derivation and the edge-on orientation of NGC 4388 would imply that NGC 4388 has a large velocity transverse to our line of sight, $V_t = (V_{\perp} - V_r \cos |i|)/\sin |i| \approx 900 \text{ km s}^{-1}$. Given the mass of the Virgo cluster, $M_{\text{Virgo}} \sim 5 - 9 \times 10^{14} \text{ M}_{\odot}$ (e.g., Tully & Shaya 1984; Giraud 1986), the total velocity of NGC 4388, $[V_t^2 + V_r^2]^{1/2} \approx 1,800 \text{ km s}^{-1}$, would imply that NGC 4388 is unbound to the Virgo cluster. We consider this unlikely based on the HI deficiency of NGC 4388 and its proximity to the center of the cluster. More likely, NGC 4388 is in a minimum potential energy part of its orbit and most of its velocity is along our line of sight. In this case, $V_{\perp} \approx V_r \cos |i| \approx 300 \text{ km s}^{-1}$, the ram pressure is 1/16th that used by Cayatte et al. (1994), and stripping occurs only in the outskirts ($R > 10 \text{ kpc}$) of NGC 4388.

However, as noted by Nulsen (1982), ram pressure stripping is not the only means by which the ICM can strip a galaxy. Momentum transfer between the hot ICM and the galaxy ISM *via* the Kelvin-Helmholtz instability can increase stripping rates under some circumstances. While the effectiveness of the ram pressure is extremely sensitive to orientation, the rate of mass loss from such turbulent viscous stripping is approximately

$$\dot{M}_{\text{visc}} = \pi R_{\text{HI}}^2 \rho_{\text{ICM}} V \approx \pi R_{\text{HI}}^2 \rho_{\text{ICM}} V_r, \quad (8)$$

where R_{HI} is the radius of the HI disk and V is the velocity of NGC 4388 relative to the cluster. Using the current extent of the HI disk of NGC 4388, Cayatte et al. (1994) estimate that $\dot{M}_{\text{visc}}(\text{NGC 4388}) \approx 1 - 6 \text{ M}_{\odot} \text{ yr}^{-1}$, implying that it will lose virtually all of its atomic content ($M_{\text{HI}} = 4 \times 10^8 \text{ M}_{\odot}$ for a distance of 16.7 Mpc; Kenney & Young 1989) during a crossing time $t_{\text{crossing}} \simeq 3 \times 10^9 \text{ yrs}$ through the cluster core. This suggests that NGC 4388 is on its first passage through the cluster core.

So far we have assumed that the ISM of NGC 4388 is experiencing the full impact of the high-velocity ICM. In reality the ICM ‘wind’ will be disturbed by the presence of the galaxy. In

particular, because the sound speed in the cluster gas is $c_s \approx 500 \text{ km s}^{-1}$ and the radial velocity of NGC 4388 relative to the Virgo cluster core is $V_r \approx 1,500 \text{ km s}^{-1}$, the interaction of the ISM of NGC 4388 with the cluster gas will be supersonic with Mach number $M = V/c_s \approx V_r/c_s \approx 3$. In this situation, a Mach cone (bow shock) forms near the leading edge of NGC 4388. The opening half-angle of the Mach cone is $\phi = 2 \tan^{-1} (1/M) \approx 40^\circ$ (note the factor of 2 missing in the expression used by Rangarajan et al. 1995). The ICM will be shocked at the Mach cone. The Rankine-Hugoniot jump conditions for adiabatic shocks ($\gamma = 5/3$) yield a post-shock velocity $V_{ps} \approx V_{\text{ICM}}/3 = 500 \text{ km s}^{-1}$, a post-shock density $n_{ps} \approx 3 n_{\text{ICM}} = 3 \times 10^{-4} \text{ cm}^{-3}$, and a post-shock temperature $T_{ps} = 3.7 T_{\text{ICM}} = 7.4 \times 10^7 \text{ K}$. From equation (1), the cooling time for the shocked ICM is $\sim 3 \times 10^{11} \text{ yrs}$, thereby justifying our assumption of an adiabatic shock. Given the rather wide opening angle of the Mach cone, the extraplanar gas observed in NGC 4388 will be affected by the shocked – rather than the pre-shock – ICM (Fig. 5).

The turbulent viscous stripping rate due to the shocked ICM will be the same as that calculated for the pre-shock ICM (because $\rho_{\text{ICM}} V_{\text{ICM}} = \rho_{ps} V_{ps}$; cf. eqn. 8), but the ram pressure on the extraplanar gas will be substantially reduced. Assuming the shocked ICM flow lines roughly follow the contours of NGC 4388 and taking $V_{\text{outflow}} = 250 \text{ km s}^{-1}$ as the velocity component of the outflowing NE cloud along the plane of the galaxy, the ram pressure from the shocked ICM on the extraplanar gas will be reduced by a factor $\rho_{\text{ICM}}/\rho_{ps} \times [(V_{\text{ICM}} - V_{\text{outflow}})/(V_{ps} - V_{\text{outflow}})]^2 \approx 8.3$ relative to that from the pre-shock ICM. Under these conditions the acceleration distance, x_{acc} , required to increase the approaching velocity of the gas in the NE cloud from 250 km s^{-1} to 350 km s^{-1} is approximately

$$x_{\text{acc}} = \frac{(V_{\text{final}}^2 - V_{\text{initial}}^2)}{2 \langle g \rangle} = \left(\frac{M_c (V_{\text{final}}^2 - V_{\text{initial}}^2)}{2C \rho_{ps} \langle (V_{ps} - V_{\text{outflow}})^2 \rangle l_c^2} \right), \quad (9)$$

where $V_{\text{final}} = 350 \text{ km s}^{-1}$, $V_{\text{initial}} = 250 \text{ km s}^{-1}$, and $\langle g \rangle$ and $\langle (V_{ps} - V_{\text{outflow}})^2 \rangle$ are time-averaged quantities. Using $M_c \approx 20 M_\odot$ and $l_c \approx 10 \text{ pc}$ for typical extraplanar line-emitting clouds [i.e., $n_c \approx 1 \text{ cm}^{-3}$ and $l_c \approx l_c(\text{HI cloud})$], we get $x_{\text{acc}} \approx 6 \text{ kpc}$. This is significantly larger than the $2 - 3 \text{ kpc}$ separation between the NE cloud and the rest of the NE complex at higher $|z|$. The implication is that the ram pressure exerted by the shocked ICM has only a small effect on the kinematics of the outflowing extraplanar material.

We conclude this section on a more speculative note by suggesting that the impact of the shocked ICM on the galaxy and the extraplanar gas component may also be cushioned by a gaseous halo in hydrostatic equilibrium around NGC 4388. As we discussed in §4.2.4, a static gaseous halo in NGC 4388 would also help to explain possible shock structures in the NE and SW clouds and the curvature of the radio jet. Our own Galaxy is known to be surrounded by a hot ($\sim 10^6 \text{ K}$) highly ionized corona extending out to several kpc from the Galactic plane (e.g., Spitzer 1990; Sembach & Savage 1992; Wang 1992). Quasar absorption-line studies (e.g., Lanzetta et al. 1995) and deep X-ray images of spiral and elliptical galaxies (e.g., Fabbiano 1988; Heckman et al. 1990; Dahlem et al. 1996) show similar gaseous halos. The main question is whether such halos survive long enough in the Virgo cluster environment. Current soft X-ray images of NGC 4388

(e.g., Matt et al. 1994; Angelo Antonelli, Matt, & Piro 1997; Iwasawa et al. 1997; Colbert et al. 1998) show soft X-ray emission of thermal origin, but the exact morphology of the X-ray gas is ill-defined because of the strong X-ray background from the Virgo ICM and the distorted halo in nearby M86 (Forman et al. 1979; White et al. 1991; Rangarajan et al. 1995). The very existence of this halo in M86, a giant E/S0 galaxy which is located close to the Virgo cluster core (15.7 Mpc according to Jacoby et al. 1990 or 18.4 Mpc according to Tonry et al. 1990) and is moving at a velocity $\sim -1,300 \text{ km s}^{-1}$ relative to the cluster ($V_r = -227 \text{ km s}^{-1}$; Jacoby et al. 1990), gives us some confidence that at least some galaxies in Virgo [albeit those more massive than NGC 4388: $M(\text{M86}) \approx 10^{12} M_\odot$; e.g., Forman et al. 1979] can keep an X-ray halo while experiencing virtually the same ram pressure as that on NGC 4388.

5. Conclusions

The complete two-dimensional coverage of our Fabry-Perot spectra on the line-emitting gas in NGC 4388 has allowed us to separate kinematically the extraplanar and disk gas components. We find that the velocity field of the high- $|z|$ gas is best described by a bipolar outflow rather than ram-pressure stripping by the fast-moving ICM in the Virgo cluster. We argue for a Mach cone with opening angle $\sim 80^\circ$ to explain these results. Unfortunately, this Mach cone will be difficult to detect, even with AXAF and XMM, because it is predicted to be roughly face-on and the strong X-ray background from nearby galaxies and the ICM in the Virgo cluster will dominate the emission. Nevertheless, high-resolution X-ray spectroscopy would be useful to constrain the velocity field (and metallicity) of the X-ray gas and to help establish if the hot gas near NGC 4388 is in hydrostatic equilibrium in the galaxy potential or instead represents fast-moving, shocked ICM. Detailed investigations of bow shocks in galaxy clusters have the potential to help us determine the space velocities of galaxy members with respect to the ICM.

From a theoretical viewpoint there is a need to explore the effects of a fast-moving ICM as it interacts with a gas-rich galaxy. Few such models have been published (e.g., Kritsuk 1984; Tosa 1994), and none considers the multi-phase nature of the ISM and the effects that this interaction may have on star formation (e.g., through compression; Kenney & Young 1989; Cayatte et al. 1994). The ICM ‘wind’ will also affect the morphology of the nuclear ejecta from starburst galaxies and AGN (e.g., wide-angle tailed radio sources in cluster environment; Loken et al. 1995). Outflowing gas will stretch in the direction of the ICM motion, thereby increasing the effective gaseous cross-section of these galaxies and perhaps contributing significantly to the population of C IV absorption lines systems in quasars (e.g., Norman et al. 1996).

We would like to thank Jim Stone for helpful discussions. SV is grateful for support of this research by a Cottrell Scholarship awarded by the Research Corporation, NASA/LTSA grant NAG 56547, and Hubble fellowship HF-1039.01-92A awarded by the Space Telescope Science Institute which is operated by the AURA, Inc. for NASA under contract No. NAS5-26555. JBH acknowledges partial support from the Fullam award of the Dudley Observatory.

REFERENCES

- Angelo Antonelli, L., Matt, G., & Piro, L. 1997, *A&A*, 317, 686
- Athanassoula, E. 1992, *MNRAS*, 259, 345
- Ayani, K., & Iye, M. 1989, *AJ*, 97, 687
- Binggeli, B., Tammann, G. A., & Sandage, A. 1987, *AJ*, 94, 251
- Bland, J., & Tully, R. B. 1988, *Nat*, 334, 43
- Blandford, R. D., & Königl, A. 1979a, *ApJ*, 232, 34
- Blandford, R. D., & Königl, A. 1979b, *Astrophys Letters*, 20, 15
- Bland-Hawthorn, J. 1995, in *Trimensional Spectroscopic Methods in Astrophysics*, IAU Coll. 149, ASP Conf. Ser. Vol. 71, eds. G. Comte and M. Marcellin, p. 72
- Böhringer, H., Briel, U. G., Schwarz, R. A., Voges, W., Hartner, G., & Trümper 1994, *Nature*, 368, 828
- Burstein, D., & Heiles, C. 1978, *ApJ*, 225, 40
- Burstein, D., & Heiles, C. 1984, *ApJS*, 54, 33
- Carral, P., Turner, J. L., & Ho, P. T. P. 1990, *ApJ*, 362, 434
- Cayatte, V., Kotanyi, C., Balkowski, C., & van Gorkom, J. H. 1994, *AJ*, 107, 1003
- Cayatte, V., van Gorkom, J. H., Balkowski, C., & Kotanyi, C. 1990, *AJ*, 100, 604
- Cecil, G. N., Bland, J., & Tully, R. B. 1990, *ApJ*, 329, 38
- Chamaraux, P., Balkowski, C., Gérard, E. 1980, *A&A*, 83, 5
- Chincarini, G., & De Souza, R. 1985, *A&A*, 153, 218
- Colbert, E. J. M., Baum, S., Gallimore, J. F., O’Dea, C. P., & Christensen, J. A. 1996a, *ApJ*, 467, 551
- Colbert, E. J. M., et al. 1996b, *ApJS*, 105, 75
- Colbert, E. J. M., Baum, S. A., O’Dea, C. P., & Veilleux, S. 1998, *ApJ*, 496, 786
- Colina, L. 1992, *ApJ*, 386, 59
- Colina, L., Fricke, K. J., Kollatschny, W., & Perryman, M. A. C. 1987, *A&A*, 186, 39
- Corbin, M. R., Baldwin, J. A., & Wilson, A. S. 1988, *ApJ*, 334, 584 (CBW)
- Cowie, L. L., & McKee, C. F. 1977, *ApJ*, 211, 135
- de Vaucouleurs, G., de Vaucouleurs, A., H. G. Jr. Corwin, Buta, R. J., Paturel, G., & Fouqué, P. 1991, *Third Reference Catalogue of Bright Galaxies* (New York, Springer-Verlag)
- DeYoung & Axford 1969, *Nature*, 216, 129
- Evans, I. N., & Dopita, M. A. 1985, *ApJS*, 58, 125

- Fabbiano, G., Kim, D.-W., & Trinchieri, G. 1992, *ApJS*, 80, 531
- Faber, S. M., et al. 1989, *ApJS*, 69, 763
- Fabricant, D., & Gorenstein, P. 1983, *ApJ*, 267, 535
- Falcke, H., Wilson, A. S., & Simpson, C. 1998, *ApJ*, 502, 199
- Fiedler, R. & Henriksen, R. N. 1984, *ApJ*, 281, 554.
- Ford, W. K., Rubin, V. C., & Roberts, M. S. 1971, *AJ*, 76, 22
- Forman, W., Schwarz, J., Jones, D., Liller, W., & Fabian, A. C. 1979, *ApJ*, 234, L27
- Gallimore, J. F., Baum, S. A., O’Dea, C. P., & Pedlar, A. 1996a, *ApJ*, 458, 136
- Gallimore, J. F., Baum, S. A., & O’Dea, C. P. 1996b, *ApJ*, 464, 198
- Giraud, E. 1986, *A&A*, 167, 25
- Goodrich, R. W., Veilleux, S., & Hill, G. 1994, *ApJ*, 433, 48
- Guhathakurta, P., van Gorkom, J. H., Kotanyi, C. G., & Balkowski, C. 1988, *AJ*, 96, 851
- Gunn, J. E., & Gott, J. R. 1972, *ApJ*, 176, 1
- Hanson, C. G., Skinner, G. K., Eyles, C. J., & Wilmore, A. P. 1990, *MNRAS*, 242, 262
- Haynes, M. P., & Giovanelli, R. 1986, *ApJ*, 306, 466
- Heckman, T. M., Miley, G. K., & Green, R. F. 1984, *ApJ*, 281, 525
- Heckman, T. M., van Bruegel, W., Miley, G. K., & Butcher, H. R. 1983, *AJ*, 88, 1077
- Hélou, G., Salpeter, E. E., & Krumm, N. 1979, *ApJ*, 228, L1
- Hélou, G., Giovanardi, G., Salpeter, E. E., & Krumm, W. 1981, *ApJS*, 46, 267
- Hélou, G., Salpeter, E. E., & Krumm, N. 1979, *ApJ*, 228, L1
- Henriksen, R. N., Vallée, J. P., & Bridle, A. H. 1981, *ApJ*, 249, 40
- Hummel, E., & Saikia, D. J. 1991, *A&A*, 249, 43
- Iwasawa, K., et al. 1997, *MNRAS*, 285, 683
- Iye, M., & Ulrich, M.-H. 1986, *Astrophysics and Space Science*, 118, 523
- Jacoby, G. H., Ciardullo, R., & Ford, H. C. 1990, *ApJ*, 356, 332
- Kenney, J. D., & Young, J. S. 1986, *ApJ*, 301, L13
- Kenney, J. D., & Young, J. S. 1989, *ApJ*, 344, 171
- Kim, D.-W., Fabbiano, G., & Trinchieri, G. 1992, *ApJS*, 80, 645
- Kinney, A. L., Antonucci, R. R. J., Ward, M. J., Wilson, A. S., & Whittle, M. 1991, *ApJ*, 377, 100
- Kukula, M. J., Pedlar, A., Baum, S. A., & O’Dea, C. P. 1995, *MNRAS*, 276, 1262
- Kritsuk, A. G. 1984, *Astrophysics*, 19, 263
- Lehnert, M. D., & Heckman, T. M. 1996, *ApJ*, 462, 651

- Lin, D. N. C., Jones, B. F., & Klemola, A. R. 1995, *ApJ*, 439, 652
- Loken, C., Roettiger, K., Burns, J. O., & Norman, M. 1995, *ApJ*, 445, 80
- Longair, M. S., Ryle, M., & Scheuer, P. A. G. 1973, *MNRAS*, 164, 243
- Matt, G., Piro, L., Antonelli, L. A., Fink, H. H., Meurs, E. J. A., & Perola, G. C. 1994, *A&A*, 292, L13
- McCall, M. L. 1984, *MNRAS*, 208, 253
- McCall, M. L., Rybski, P. M., & Shields, G. A. 1985, *ApJS*, 57, 1
- McKee, C. F., & Cowie, L. L. 1975, *ApJ*, 195, 715
- McKee, C. F., Cowie, L. L., & Ostriker, J. P. 1978, *ApJ*, 219, L23
- Morse, J. A., Cecil, G., Wilson, A. S., & Tsvetanov, Z. I. 1998, *ApJ*, 505, 159
- Mould, J., Aaronson, M., & Huchra, J. 1980, *ApJ*, 238, 458
- Mulchaey, J. S., Wilson, A. S., & Tsvetanov, Z. 1996, *ApJ*, 467, 197
- Norman, C. A., Bowen, D. V., Heckman, T., Blades, C., & Danly, L. 1996, *ApJ*, 472, 73
- Nulsen, P. E. J. 1982, *MNRAS*, 198, 1007
- O'Donoghue, A. A., Eilek, J. A., & Owen, F. N. 1993, *ApJ*, 408, 428
- Osterbrock, D. E., & Martel, A. 1992, *PASP*, 104, 76
- Osterbrock, D. E., Tran, H. D., & Veilleux, S. 1992, *ApJ*, 389, 196
- Pedlar, A., Dyson, J. E., & Unger, S. W. 1985, *MNRAS*, 214, 463
- Petitjean, P., & Durret, F. 1993, *A&A*, 277, 365
- Phillips, M. M., & Malin, D. F. 1982, *MNRAS*, 199, 205
- Piner, B. G., Stone, J. M., & Teuben, P. J. 1995, *ApJ*, 449, 508
- Pierce, M. J., & Tully, R. B. 1988, *ApJ*, 330, 579
- Pogge, R. W. 1988, *ApJ*, 332, 702
- Prieto, M. A., & Freudling, W. 1996, *MNRAS*, 279, 63
- Rangarajan, F. V. N., White, D. A., Ebeling, H., & Fabian, A. C. 1995, *MNRAS*, 277, 1047
- Rubin, V. C., Kenney, J. D. P., & Young, J. S. 1997, *AJ*, 113, 1250
- Sandage, A. 1978, *AJ*, 83, 904
- Sandage, A., & Tammann, G. 1984, *Nature*, 307, 326
- Sembach, K. R., & Savage, B. D. 1992, *ApJS*, 83, 147
- Sgro, A. G., 1975, *ApJ*, 197, 621
- Shields, J. C., & Filippenko, A. V. 1988, *ApJ*, 332, L55
- Shields, J. C., & Filippenko, A. V. 1996, *A&A*, 311, 393

- Shopbell, P. L., & Bland-Hawthorn, J. 1998, *ApJ*, 493, 129
- Smith, M. D., & Norman, C. A. 1981, *MNRAS*, 194, 771
- Spitzer, L. 1990, *ARAA*, 28, 71
- Stone, J. L., Wilson, A. S., & Ward, M. J. 1988, *ApJ*, 330, 105
- Storchi-Bergmann, T., Bica, E., & Pastoriza, M. G. 1990, *MNRAS*, 245, 749
- Takano, S., & Koyama, K., 1991, *PASJ*, 43, 1
- Taylor, D., Dyson, J. E., Axon, D. J., & Pedlar, A. 1989, *MNRAS*, 240, 487
- Tosa, M. 1994, *ApJ*, 426, L81
- Tonry, J. L., Ajhar, E. A., & Luppino, G. A. 1990, *AJ*, 100, 1416
- Tsvetanov, Z.I., & Walsh, J.R. 1992, *ApJ*, 386, 485
- Tully, R. B., & Shaya, E. J. 1984, *ApJ*, 281, 31
- Veilleux, S., & Bland-Hawthorn, J. 1997, *ApJ*, 479, L105
- Veilleux, S., Bland-Hawthorn, J., & Cecil, G. 1995, *ApJ*, 445, 152
- Veilleux, S., Bland-Hawthorn, J., & Cecil, G. 1999, *ApJ*, submitted (VBC)
- Veilleux, S., Cecil, G., Bland-Hawthorn, J., Tully, R. B., Filippenko, A. V., & Sargent, W. L. W. 1994, *ApJ*, 433, 48
- Veilleux, S., Goodrich, R. W., & Hill, G. J. 1997, *ApJ*, 477, 631
- Veilleux, S., Tully, R. B., & Bland-Hawthorn, J. 1993, *AJ*, 105, 1318
- Walker, I. R., Mihos, J. C., & Hernquist, L. 1996, *ApJ*, 460, 121
- Wang, Q. D. 1992, *PASP*, 104, 812
- Wehrle, A. E., & Morris, M. 1987, *ApJ*, 313, L43
- White, D. A., Fabian, A. C., Forman, W., Jones, C., & Stern, C. 1991, *ApJ*, 375, 35
- Whittle, M., Pedlar, A., Meurs, E. J. A., Unger, S. W., Axon, D. J., & Ward, M. J. 1988, *ApJ*, 326, 125
- Wilson, A. S. 1981, in *Proc. Second ESO/ESA Workshop on Optical Jets in Galaxies*, ed. A. F. M. Moorwood & K. Kj  r (Garching: ESO), 125
- Wilson, A. S., & Tsvetanov, Z. 1994, *AJ*, 107, 1227
- Woods, D., Madore, B. F., & Fahlman, G. G. 1990, *ApJ*, 353, 90
- Yasuda, N., Fukugita, M., & Okamura, S. 1997, *ApJS*, 108, 417

Table 1: Kinematical Parameters of NGC 4388^(a)

Component	Parameter	NGC 4388
Disk	V_{sys}	2,525 km s ⁻¹
	V_{max} (deprojected)	240 km s ⁻¹
	P.A. (major axis)	90°
	Inclination i	-78°
Bar	P.A. (intrinsic to disk)	135°
	Inner radius R_b	1.5 kpc
	Orbit eccentricity $R \leq R_b$	0.3
	Outer radius R_d	5.1 kpc

^(a) Refer to Veilleux, Bland-Hawthorn, & Cecil (1999) for a detailed discussion of these parameters.

Fig. 1.— The distributions of the line emission in NGC 4388. (top) $H\alpha$; (bottom) $[O\ III]\ \lambda 5007$. The optical fluxes were determined from Gaussian fitting of the emission-line profiles. North is at the top and East to the left. The spatial scale, indicated by a horizontal bar at the bottom of the image, corresponds to $12''$, or 1 kpc for the adopted distance of 16.7 Mpc. The optical continuum nucleus is indicated in each panel by a cross. The flux scale is logarithmic and is expressed in units of $10^{-16}\ \text{erg s}^{-1}\ \text{cm}^{-2}\ \text{arcsec}^{-2}$.

Fig. 2.— $[O\ III]\ \lambda 5007/H\alpha$ flux ratio map of the line-emitting gas in NGC 4388. Same spatial scale and orientation as Fig. 1. The optical continuum nucleus is indicated by a cross. Values for this ratio range from ~ 0.1 in the disk to $\gtrsim 5$ in the extraplanar material. Regions with $[O\ III]\ \lambda 5007/H\alpha = 9$ indicate $[O\ III]$ emission clouds where no $H\alpha$ was detected.

Fig. 3.— Barycentric velocity field of the line-emitting gas in NGC 4388. (top) $H\alpha$; (bottom) $[O\ III]\ \lambda 5007$. Same spatial scale and orientation as Fig. 1. The optical continuum nucleus is indicated in each panel by a cross. The velocities range from about $-200\ \text{km s}^{-1}$ to $+200\ \text{km s}^{-1}$ relative to systemic ($= 2,525\ \text{km s}^{-1}$).

Fig. 4.— Models of the disk velocity field: (top left) a simple axisymmetric kinematic model, (top right) a kinematic model with elliptical streaming motion in the center. See text and Table 1 for more details on the parameters of the models. The residuals after subtracting the axisymmetric model from the observed $H\alpha$ velocity field are presented in the bottom left panel while the bottom right panel shows the residuals after subtracting the non-axisymmetric model. The non-axisymmetric model is clearly a better representation of the data.

Fig. 5.— Geometry of the ICM – NGC 4388 interaction. The values of the parameters indicated on this figure and discussed in the text are: $i = -78^\circ$, $\phi = 40^\circ$, $\theta > 12^\circ$, $R_{H\ I} = 10\ \text{kpc}$, $V_{\text{ICM}} = 1,500\ \text{km s}^{-1}$, $n_{\text{ICM}} \sim 10^{-4}\ \text{cm}^{-3}$, $V_{\text{ps}} = 500\ \text{km s}^{-1}$, $n_{\text{ps}} \sim 3 \times 10^{-3}\ \text{cm}^{-3}$. The observer is located on the right in the same plane as the figure and at a distance of 16.7 Mpc.




Fig. 1.—




Fig. 2.—




Fig. 3.—




Fig. 4.—




Fig. 5.—